

**Reference list**  
**IMAP**  
**Fishes**

- 1) Angermeier, P.L., and Karr, J.R. 1986. *Applying and index of biotic integrity based on stream-fish communities: considerations in sampling and interpretation*. North American Journal of Fisheries Management. **6**: 418-429.
- 2) Angermeier, P.L., Smogor, R.A., and Steele, S.D. 1991. *An electric seine for collecting fish in streams*. North American Journal of Fisheries Management. **11**:352-357.
- 3) Angermeier, P.L., and R.A. Smogor. 1995. *Estimating number of species and relative abundances in stream-fish communities: effects of sampling effort and discontinuous spatial distributions*. Can. J. Fish. Aquat. Sci. **52**:936-949.
- 4) Bain, M. B., and Stevenson, N. J. Editors. 1999. *Aquatic habitat assessment: common methods*. American Fisheries Society, Bethesda, Maryland.
- 5) Bayley, P.B., and Herendeen, R.A. 2000. *The efficiency of a seine net*. Trans. Am. Fish. Soc. **129**:901-923.
- 6) Benson, J.B., and Magnuson, J.J. 1992. *Spatial heterogeneity of littoral fish assemblages in lakes: relation to species diversity and habitat structure*. Can. J. Fish. Aquat. Sci. **49**: 1493-1500.
- 7) Bohlin, T., and Sundstrom, B. 1997. *Influence of unequal catchability on population estimates using the Lincoln index and the removal method applied to electro-fishing*. Oikos. **28**:123-129.
- 8) Bosley, T.R., Pennington, C.H., Potter, M.E., and Knight, S.S. 1986. *Evaluation of larval fish sampling gears for use on large rivers*. Technical Report E-86-9, U.S. Army Engineers Waterways Experiment Station, Vicksburg, Miss.
- 9) Bros, W.E., Cowell, B.C. 1987. *A technique for optimizing sample size (replication)*. Journal of Experimental Marine Biology and Ecology. **114**: 63-71.
- 10) Cole, R.A., MacMillan, J.R. 1984. *Sampling larval fish in the littoral zone of western Lake Erie*. J. Great Lakes Res. **10**(1): 15-27.
- 11) Connolly, R.M. 1999. *Saltmarsh as habitat for fish and nektonic crustaceans: challenges in sampling designs and methods*. Australian J. of Ecology. **24**:422-430.
- 12) Dewey, M. R., Holland-Bartels, L.E., and Zigler, S.J. 1989. *Comparison of fish catches with buoyant pop nets and seines in vegetated and non-vegetated habitats*. N. Am. J. Fish. Management. **9**:249-253.
- 13) Dewey, M.R. 1992. *Effectiveness of a drop net, a pop net, and an electrofishing frame for collecting quantitative samples of juvenile fishes in vegetation*. North American Journal of Fisheries Management. **12**:808-813.
- 14) Dixon, P.M., and Garrett, K.A. 1993. *Sampling ecological information: choice of sample size, reconsidered*. Ecological Modeling. **68**:67-73.
- 15) Eckblad, J.W. 1991. *How many samples should be taken?* Bioscience. **41**(5): 346-348.
- 16) Espegren, G.D., and Bergesen, E.P. 1990. *Quantitative sampling of fish populations with a mobile rising net*. North American Journal of Fisheries Management. **10**:469-478.
- 17) Fairweather, P.G. 1991. *Statistical power and design requirements for environmental monitoring*. Aust. J. Freshwater Res. **42**:555-567.
- 18) Fitzpatrick, F. A., Waite, I. R., D'Arconte, P. J., Meador, M. R., Maupin, M. A., and Gurtz, M. E. 1998. *Revised methods for characterizing stream habitat in the national water quality assessment program*. Water Resources Investigations Report 98-4052. United States Geological Survey. Raleigh, North Carolina
- 19) Flosi, G. S., Downie, J., Hopelain, M., Bird, B., and Collins. 1998. *California salmonid stream habitat restoration manual, 3<sup>rd</sup> edition*. CA Resources Agency, Department of Fish and Game.
- 20) Franson, M. A. H. Managing editor. 1989. *Standard Methods for the Examination of Water and Wastewater, 17 ed*. American Public Health Association. Washington D.C.
- 21) Gardiner, W.R. 1984. *Estimating population densities of salmonids in deep water in streams*. J. Fish Bio. **24**:41-49.
- 22) Gelwick, F.P. 1990. *Longitudinal and temporal comparisons of riffle and pool fish assemblages in a northeastern Oklahoma Ozark stream*. Copeia. **4**:1072-1082.
- 23) Gorman, O.T., J.R. Karr. 1978. *Habitat Structure and Stream Fish Communities*. Ecology. **59**(3):507-515.
- 24) Green, R. H., and Young, R.C. 1993. *Sampling to detect rare species*. Ecological Applications. **3**(2): 351-356.
- 25) Hankin, D.G. 1984. *Multistage sampling design in fisheries research: applications in small streams*. Can. J. Fish. Aquat. Sci. **41**:1575-1591.
- 26) Hankin, D.G., and Reeves, G.H. 1988. *Estimating total fish abundance and total habitat area in small streams based on visual estimation methods*. Can. J. Fish. Aquat. Sci. **45**:834-844.
- 27) Hanson, J.M., and Leggett, W.C. 1982. *Empirical prediction of fish biomass and yield*. Can. J. Fish. Aquat. Sci. **39**:257-263.
- 28) Harris, J.M. 1995. *The use of fish in ecological assessments*. Australian Journal of Ecology. **20**: 65-80.
- 29) Heggenes, J., Brabrand, A., and Saltveit, S.J. 1990. *Comparison of three methods for studies of stream habitat use by young brown trout and Atlantic salmon*. Transactions of the American Fisheries Society. **119**:101-111.
- 30) Heimbuch, D.G., Wilson, H.T., Weisberg, S.B., Volstad, J.H., and Kazyak, P. F. 1997. *Estimating fish abundance in stream surveys by using double-pass removal sampling*. Trans. Am. Fish. Soc. **126**:795-803.
- 31) Hewitt, D.P. 1979. *Tests to confirm quantitative sampling of young fish by the Bagenal buoyant net*. Freshwater Biology. **9**:339-341.
- 32) Hewitt, D.P., McBride, G.B., Pridmore, R.D. and Thrush, S.F. 1993. *Patchy distributions: Optimizing sample size*, Environmental Monitoring and Assessment. **27**: 95-105.
- 33) Hinch, S.G., 1991. *Small- and large-scale studies in fisheries ecology: the need for cooperation among researchers*. Fisheries. **16**(3): 22-27.
- 34) Hinch, S.G., Collins, N.C., and Harvey, H.H. 1991. *Relative abundance of littoral zone fishes: biotic interaction, abiotic factors, and postglacial colonization*. Ecology. **72**(4):1314-1324.

- 35) Hinch, S.G., Somers, K.M., and Collins, N.C. 1994. *Spatial autocorrelation and assessment of habitat-abundance relationships in littoral zone fish*. Can. J. Fish. Aquat. Sci. **51**:701-712.
- 36) Jackson, D.A., Somers, K.M., Harvey, H.H. 1992. *Null models and fish communities: evidence of nonrandom patterns*. The American Naturalist. **139**(5):930-951.
- 37) Jackson, D.A., and Harvey, H.H. 1993. *Community concordance and community-environment relationships*. Can. J. Fish. Aquat. Sci. **50**:2641-2651.
- 38) Jackson, D.A., and Harvey, H.H. 1997. *Qualitative and quantitative sampling of lake fish communities*. Can. J. Fish. Aquat. Sci. **54**:2807-2813.
- 39) Jacobsen, T., and Kushlan, J.A. 1987. *Sources of sampling bias in enclosure fish trapping: effects on estimates of density and diversity*. Fisheries Research **5**:401-412.
- 40) Kjelson, M.A., and Johnson, G.N. 1973. *Description and evaluation of a portable drop-net for sampling nekton populations*. Proceedings of the Southeast Association of Game and Fish Commissioners. **27**: 653-662.
- 41) Kjelson, M.A., and Johnson, G.N. 1978. *Catch efficiency of a 6.1m trawl for estuarine fish populations*. Transactions of the American Fisheries Society. **107**: 246-254.
- 42) Kjelson, M.A., Turner, W.R., Johnson, G.N. 1975. *Description of a stationary drop-net for estimating nekton abundance in shallow waters*. Transactions of the American Fisheries Society. **1**:46-49.
- 43) Kruse, C.G., Hubert, W.A., and Rahel, F.J. 1998. *Single-pass electrofishing predicts trout abundance in mountain streams with sparse habitat*. North American Journal of Fisheries Management. **18**: 940-946.
- 44) Kushlan, J.A. 1974. *Quantitative sampling of fish populations in shallow, freshwater environments*. Transactions of the American Fisheries Society. **2**:348-352.
- 45) Kushlan, J.A., 1981. *Sampling characteristics of enclosure fish traps*. Transactions of the American Fisheries Society. **110**: 557-562.
- 46) Lai, H. 1993. *Optimal sampling design for using the age-length key to estimate age composition of a fish population*. Fishery Bulletin. **92**: 382-388.
- 47) Larson, E.W., Johnson, D.L., and Lynch W.E., Jr., 1986. *A buoyant pop net for accurately sampling fish at artificial habitat structures*. Transactions of the American Fisheries Society. **115**:351-355.
- 48) Lobon-Cervia, J., and Utrilla, C.G. 1993. *A simple model to determine stream trout (Salmo trutta L.) densities based on one removal with electrofishing*. Fisheries Research. **15**: 369-378.
- 49) Loesch, J., Bishop, H., Crowe, A., Huckyr, R., Wagner, P. 1976. *Technique for estimating trawl efficiency in catching brown shrimp (Panaeus aztecus), Atlantic croaker (Micropogon undulates), and spot (Leiostomus xanthurus)*. Gulf Research Reports. **5**: 29-33.
- 50) Lyons, J. 1986. *Capture efficiency of a beach seine for seven freshwater fishes in a north-temperate lake*. North American Journal of Fisheries Research. **6**:288-289.
- 51) Lyons, J. 1992. *The length of stream to sample with a towed electrofishing unit when fish species richness is estimated*. North American Journal of Fisheries Management. **12**: 198-203.
- 52) Mahon, R. 1980. *Accuracy of catch-effort methods for estimating fish density and biomass in streams*. Biol. Fish. **5**(4): 343-360.
- 53) Manly, B.F.J. 1992. *Bootstrapping for determining sample sizes in biological studies*. J. Exp. Biol. Ecol. **158**: 189-196.
- 54) Minns, C.K., Cairns, V.W., Randall, R.G., and Moore, J.E. 1994. *An index of biotic integrity (IBI) for fish assemblages in the littoral zone of Great Lakes' areas of concern*. Can. J. Fish. Aquat. Sci. **51**: 1804-1822.
- 55) Mitro, M.G., and Zale, A.V. 2000. *Predicting fish abundance using single-pass removal sampling*. Can. J. Fish. Aquat. Sci. **57**:951-961.
- 56) Moyle, P. B., Yoshiyama, R. M., Williams, J. E., and Wikramanayaki, E. D. 1995. *Fish species of special concern in California 2<sup>nd</sup> edition*. University of California, Davis.
- 57) Nielson, L.A., Johnson, D.L and Editors. 1983. *Fisheries techniques*. American Fisheries Society, Bethesda, Maryland.
- 58) Nielsen, J.L. 1998. *Electrofishing California's endangered fish populations*. Fisheries. **23**(12): 6-12.
- 59) Northcote, T.G., and Wilkie, D.W. 1963. *Underwater census of stream fish populations*. Transactions of the American Fisheries Society. **92**:146-151.
- 60) Palmer, M.W., and White, P.S. 1994. *Scale dependence and the species area relationship*. The American Naturalist. **144**(5): 717-740.
- 61) Parsley, M.J., Palmer, D.E., and Burkhardt, R.W. 1989. *Variation in capture efficiency of a beach seine for small fishes*. North American Journal of Fisheries Management. **9**:239-244.
- 62) Peterman, R.M. 1990. *Statistical power analysis can improve fisheries research and management*. Can. J. Fish. Aquat. Sci. **47**:2-15.
- 63) Peterson, N.P., and Cederholm, C.J. 1984. *A Comparison of the removal and mark-recapture methods of population estimation for juvenile coho salmon in small streams*. North American Journal of Fisheries Management. **4**:99-102.
- 64) Pierce, C.L., Rasmussen, J.B., and Leggett, W.C. 1990. *Sampling littoral fish with a seine: corrections for variable capture efficiency*. Can. J. Fish. Aquat. Sci. **47**:1004-1010.
- 65) Pollock, K.H. 1982. *A capture-recapture design robust to unequal probability of capture*. J. Wildl. Manage. **46**(3): 752-756.
- 66) Pringle, J.D. 1984. *Efficiency estimates for various quadrat sizes used in benthic sampling*. Can. J. Fish. Aquat. Sci. **41**:1485-1489.
- 67) Pusey, B.J., Kennard, M.J., Arthur, J.M., and Arthington. A.H. 1998. *Quantitative sampling of stream fish assemblages: single- vs multiple-pass electrofishing*. Australian Journal of Ecology. **23**:365-374.
- 68) Rahel, F.J. 1984. *Factors structuring fish assemblages along a bog lake successional gradient*. Ecology. **65**(4):1276-1289.
- 69) Riley, S.R., and Fausch, K.D. 1992. *Underestimation of trout population size by maximum-likelihood removal estimates in small streams*. Transactions of the American Fisheries Society. **12**:768-776.
- 70) Rossier, O. 1997. *Comparison of gillnet sampling and night visual census of fish communities in the littoral zone of Lake Geneva, Switzerland*. Arch. Hydrobiol. **139**(2): 223-233.
- 71) Rozas, L.P. 1992. *Bottomless lift net for quantitatively sampling nekton on intertidal marshes*. Marine Ecology Progress Series. **89**:287-292.
- 72) Rozas, L.P., and Minello, T.J. 1997. *Estimating densities of small fishes and decapod crustaceans in shallow estuarine habitats: a review of sampling design with focus on gear selection*. Estuaries. **20**(1): 199-213.
- 73) Rundle, H.D., and Jackson, D.A. 1996. *Spatial and temporal variation in littoral-zone fish communities: a new statistical approach*. Can. J. Fish. Aquat. Sci. **53**:2167-2176.
- 74) Ryder, R.A. 1965. *A method for estimating potential fish production of north-temperate lakes*. Trans. Am. Fish. Soc. **94**: 214-218.
- 75) Seber, G.A.F., and Le Cren, E.D. *Estimating population parameters from catches large relative to the population*. 631-643.
- 76) Serafy, J.E., R.M. Harrell, J.C. Stevenson. 1988. *Quantitative sampling of small fishes in dense vegetation: design and field testing of portable "pop-nets"*. J. Appl. Ichthyol. **4**:149-157.

- 77) Strange, C.D., Aprahamian, M.W., and Winstone A.J. 1989. *Assessment of a semi-quantitative electric fishing sampling technique for juvenile Atlantic salmon, Salmo salar L., and trout, Salmo trutta L., in small streams.* Aquaculture and Fisheries Management. **20**:485-492.
- 78) Sullivan, L.F., Emiliani, D.A., and Baxter, K.N. 1985. *Standing stock of juvenile brown trout, Panaeus azteca, in Texas coastal ponds.* Fishery Bulletin. **83**: 677-682.
- 79) Taylor, C.M. 1996. *Abundance and distribution within a guild of stream fishes: local processes and regional patterns.* Freshwater Biology. **36**:385-396.
- 80) Taylor, C.M. 1997. *Fish species richness and incidence patterns in isolated and connected stream pools: effects of pool volume and spatial position.* Oecologia. **110**:560-566.
- 81) Taylor, C.M. 2000. *A large-scale comparative analysis of riffle and pool fish communities in an upland stream system.* Environmental Biology of Fishes. **58**:89-95.
- 82) Tonn, W.M., and Magnuson, J.J. 1982. *Patterns in the species composition and richness of fish assemblages in northern Wisconsin lakes.* Ecology. **63**(4): 1149-1166.
- 83) Weaver, M.J., Magnuson, J.J., and Clayton, M.K. 1993. *Analyses for differentiating littoral fish assemblages with catch data from multiple sampling gears.* Trans. Am. Fish. Soc. 122:1111-1119.
- 84) Wiley, M.L., and Tsai, C. 1983. *The relative efficiencies of electrofishing vs. seines in Piedmont streams of Maryland.* North American Journal of Fisheries Management. **3**:243-253.
- 85) Zalewski, M. 1983. *The influence of fish community structure on the efficiency of electrofishing.* Fish. Mgmt. **14**(4): 177-186.
- 86) Zedler, J.B., and Langis, R. 1990. *A manual for assessing restored and natural coastal wetlands with examples from southern California.* Pacific Estuarine Research Laboratory. California State University, San Diego.
- 87) Zippin, C. 1956. *An evaluation of the removal method of estimating animal populations.* Biometrics. **12**: 163-169.